INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) (51) International Patent Classification 6: (11) International Publication Number: WO 95/11671 A61K 9/14, 9/00, 9/20 A1 (43) International Publication Date: 4 May 1995 (04.05.95) (21) International Application Number: (81) Designated States: AU, CA, CN, JP, PL, RU, European patent PCT/US94/12018 (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, (22) International Filing Date: 20 October 1994 (20.10.94) NL, PT, SE). (30) Priority Data: Published 08/144,592 28 October 1993 (28,10,93) US With international search report. (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: BRIDEAU, Michelle, Elizabeth; Apartment G, 8428 Jonathan Lane, Maineville, OH 45039 (US). CARELLA, Anne, Marie; 2339 Park Avenue, Cincinnati, OH 45206 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US). (54) Title: FAST DISSOLVING DOSAGE FORMS CONTAINING MAGNESIUM ALUMINUM SILICATE AND MULTIPLE ACTIVE **INGREDIENTS** (57) Abstract An adsorbate composition comprising magnesium aluminum silicate and two or more pharmaceutically acceptable actives in a fast dissolving dosage form.

÷.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

		CP	Third Visadon	MR	Mauritania
AT	Austria	GB	United Kingdom	MW	Malawi
ΑU	Australia	GE	Georgia		
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NŁ	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	Æ	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SI	Slovenia
CI	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	LI	Liechtenstein	SN	Senegal
CN	China .	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA.	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali	UZ.	Uzbekistan
FR	France	MN	Mongolia	VN	Viet Nam
GA	Gabon				

TECHNICAL FIELD

The present invention relates to a good tasting, fast dissolving adsorbate composition consisting of magnesium aluminum silicate with two or more pharmaceutically acceptable active ingredients.

BACKGROUND OF THE INVENTION

Adsorbates have been used for many years, masking the untoward taste of many compounds. The use of magnesium aluminum silicate in this capacity is disclosed in, for example, <u>U.S. Patent 4.711.774</u>, December 8, 1987; <u>4.716.033</u>, December 29, 1987; <u>4.758.424</u>, July 19,1988; <u>4.758.425</u>, July 19,1988; <u>4.761.274</u>, August 2,1988 to Denick, Jr. et al., <u>U.S. Patent 4.753,800</u>, June 28, 1988 to Mozda and <u>U.S. Patent 3,140,978</u>, July 14, 1964 to Zentner.

While the prior art discloses magnesium aluminum silicate as useful in masking the taste of pharmacologically active compounds, there is still a need for fast dissolving, taste-masking formulations of this kind incorporating multiple pharmacologic actives. The prior art disclosures of immediate release dosage forms containing magnesium aluminum silicate as the taste masking agent generally focus on masking the taste of a single pharmaceutically active ingredient. This may be attributed to the assumption that any increase in the amount of the bitter tasting component(s) used (e.g. higher doses of the initial compound or the admixing of an additional compound(s)) would require a corresponding increase in the amount of magnesium aluminum silicate used. The causal link between this assumption and the focus of the prior art becomes evident as one appreciates the problems that arise with increasing amounts of magnesium aluminum silicate. These problems include decreased disintegration time, decreased release of the pharmaceutically active ingredient and decreased palatability (i.e. experiencing an undesirable clay taste). The present inventors, however, have discovered that despite the presence of multiple. pharmaceutically active compounds, fast dissolving dosage forms which incorporate a single taste masking agent, magnesium aluminum silicate, still retain their taste masking ability even at levels low enough to avoid the above-mentioned problems. It is therefore an object of the present invention to provide a good-tasting adsorbate composition that contains two or more pharmacologic actives. It is a further object of the present invention to provide a good-tasting, fast dissolving composition incorporating said pharmaceutical actives. Still a further object is to provide a

10

15

20

25

30

35

method of making a good tasting, quick dissolving medicament containing multiple pharmaceutical actives.

These objectives and additional objectives will become readily apparent from the detailed description which follows.

SUMMARY OF THE INVENTION

The present invention relates to a pharmaceutical composition comprising:

- (a) a clay component consisting of magnesium aluminum silicate;
- (b) a safe and effective amount of two or more pharmaceutically acceptable active ingredients selected from the group consisting of: an antitussive; an antinauseant; a nutritional supplement; a laxative; an appetite suppressant; an analgesic; an antiasthmatic; an antihistamine; a decongestant; an expectorant; an antacid; an antidiarrheal, a H₂-Receptor antagonist and mixtures thereof; and
- (c) a pharmaceutically acceptable carrier wherein said pharmaceutically acceptable carrier can be rapidly disintegrated in aqueous solution.

The present invention further relates to methods of treating symptoms such as those associated with the common cold, respiratory disorders, cough, cold, cold-like and/or flu symptoms associated with the common cold, gastrointestinal disorders and allergies; comprising the administration of a safe and effective amount of the compositions of the present invention.

All percentages and ratios herein are by weight unless otherwise specified. Additionally, all measurements are made at 25°C unless otherwise specified.

By "safe and effective amount," as used herein, is an amount that is effective to mitigate and/or treat the symptoms for which the active ingredient is indicated in a human without undue adverse side effects commensurate with a reasonable risk/benefit ratio.

DETAILED DESCRIPTION OF THE INVENTION

The compositions of the present invention contain essential components as well as various nonessential components as indicated below.

ESSENTIAL COMPONENTS

Magnesium Aluminum Silicate

5

10

15

20

25

30

35

The first essential component of the present invention is a safe and effective amount of magnesium aluminum silicate of the formula Al2MgO8Si2. Magnesium aluminum silicate occurs naturally in such smectite minerals as colerainite, saponite, sapphirine, sheridanite zebedassite and has been used extensively in a variety of cos-

metic and pharmaceutical formulations. The refined magnesium aluminum silicates used in the present example, Veegum® HS, is available from and manufactured by R.T. Vanderbilt Company, Inc., Norwalk, CT. A Typical chemical analysis of Veegum® HS, is as follows:

5	Silicon dioxide	63.0
	Magnesium oxide	10.5
	Aluminum oxide	10.5
	Ferric oxide	0.9
	Calcium oxide	2.3
10	Sodium oxide	2.4
	Potassium oxide	1.2
	Ignition loss	7.5

having a combined density of 2.6mg/m².

15

20

25

30

35

The compositions of this invention typically comprise from about 0.01% to about 50%, preferably from about 0.1% to about 25%, more preferably from about 1.0% to about 10% and most preferably from about 1.0% to about 5%, by weight of a magnesium aluminum silicate.

Magnesium aluminum silicates are described more fully in, for example, <u>U.S. Patent 4.761.274</u>, August 2, 1988, to Denick, Jr. et al., <u>U.S. Patent 4.753.800</u>, June 28, 1988, to Mozda, and <u>U.S. Patent 3.140.978</u>, July 14, 1964 to Zentner; all of which are herein incorporated by reference.

Pharmaceutically acceptable actives

The two or more pharmaceutically acceptable actives useful in the present invention may be selected from among the various groups of chemical compounds or materials suitable for oral administration and having a pharmacological action. These pharmaceutically acceptable active compounds or materials should be compatible with the other essential ingredients and compatible in combination with other included active materials or compounds. The pharmaceutically acceptable active compounds or materials are present at a level from about 0.01% to about 75%, preferably from about 0.1% to about 50%, more preferably from about 1.0% to about 25% and most preferably from about 1.0% to about 10%. Useful pharmaceutically acceptable active materials or compounds may include, but are not limited to: bronchodilators, anorexiants, antihistamines, nutritional supplements (such as vitamins, minerals, fatty acids, amino acids, and the like), laxatives, analgesics, antacids, H2-receptor antagonists, antidiarrheals, decongestants, antitussives, antinauseants, antimicrobials, antifungals, antivirals, expectorants, anti-inflammatory agents, antipyretics, their pharmaceutically acceptable salts and mixtures thereof.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include sodium, potassium, lithium, ammonia, calcium, magnesium, ferrous, zinc, manganous, aluminum, ferric, manganic salts and the like. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, tertiary and quaternary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as triethylamine, tripropylamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, lysine, arginine, histidine, caffeine, procaine, N-ethylpiperidine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglycamine, theobromine, purines, piperazine, piperidine, polyamine resins and the like.

Examples of decongestants useful in the compositions of the present invention include pseudoephedrine, phenylpropanolamine, phenylephrine and ephedrine, their pharmaceutically acceptable salts, and mixtures thereof.

Examples of antitussives useful in the compositions of the present invention include dextromethorphan, chlopedianol, carbetapentane, caramiphen, noscapine, diphenhydramine, codeine, hydrocodone, hydromorphone, their pharmaceutically acceptable salts, and mixtures thereof.

Examples of expectorants (also known as mucolytic agents) useful in the present invention include; glyceryl guaiacolate, terpin hydrate, ammonium chloride, Nacetylcysteine, and ambroxol, their pharmaceutically acceptable salts, and mixtures thereof.

Examples of analgesics useful in the present invention include; morphine, codeine, meperidine, pentazocine, propoxyphene, acetaminophen, allopurinol, acetylsalicylic acid, choline salicylate, ketoprofen, magnesium silicate, fenoprofen, ibuprofen, indomethacin, naproxen, and many others and their pharmaceutically acceptable salts and mixtures thereof.

Examples of antihistamines useful in the present invention include; brompheniramine, chlorpheniramine, clemastine, dexchlorpheniramine, diphenhydramine, doxylamine, promethazine, terfenadine, triprolidine and many others and their pharmaceutically acceptable salts and mixtures thereof.

Analgesics, decongestants, antihistamines, expectorants and antitussives, as well as their acceptable dosage ranges are described in <u>U.S. Patent 4,783,465</u> to Sunshine et al., issued November 8, 1988, and <u>U.S. Patent 4,619,934</u> to Sunshine et al., issued October 28, 1986, which are incorporated by reference herein.

Examples of gastrointestinal agents suitable for use in the present invention include anticholinergies, including atropine, clidinium and dicyclomine; antacids, in-

15

10

20

25

30

cluding aluminum hydroxide, bismuth subsalicylate, simethicone, calcium carbonate and magaldrate; H2-receptor antagonists including: cimetidine, famotidine, nizatidine and ranitidine; laxatives, including: phenolphthalein and casanthrol; and antidiarrheals including: diphenoxylate and loperamide.

5

10

15

20

25

35

Further examples of suitable analgesics, decongestants, antitussives, expectorants and antihistamines as well as bronchodilators, anorexiants, laxatives, antiemetics, antimicrobials, antibacterials, antifungals, anti-inflammatory agents, antivirals, antipyretics, nutritional supplements, anticholinergics, antacids, H2-receptor antagonists, antidiarrheals and other miscellaneous gastrointestinal compounds and their acceptable dosage ranges are described in Remington's Pharmaceutical Sciences, pp. 734-789, 791-799, 861-868, 907-945, 875-888, 1002-1034, 1098-1121, 1124-1131, 1173-1224, 1232-1241, (Alfonso R.Gennaro, editor) (18th ed. 1990), herein incorporated by reference.

NONESSENTIAL COMPONENTS

Persons skilled in the art will quickly realize many other ingredients will be suitable for inclusion into the present invention. Nonessential components include, but are not limited to: coloring agents; flavoring agents, including: vanilla, cherry, grape, orange, peppermint, spearmint, anise, blueberry raspberry, banana, chocolate, caramel, strawberry, lemon, menthol and ProsweetTM MM50 (a combination of natural and artificial flavors and propylene glycol, available from Virginia Dare Extract Co., Inc., Brooklyn, NY); sweeteners, including saccharin, dextrose, levulose, sucrose, cyclamate, mannitol and aspartate, along with many others; suspending agents, including xanthum gum, acacia gum, carboxymethylcellulose, starch and methylcellulose; preservatives; releasing agents, including polysorbate 80, sodium lauryl sulfate, vegetable oils and magnesium stearate; and water.

Another preferred nonessential component of the present invention is a cooling agent or a combination of cooling agents. Suitable cooling agents are those described in <u>U.S. Patent 4,136,163</u>, January 23, 1979, to Watson et al., <u>U.S. Patent 4,230,668</u>, October 28, 1980, to Rowsell et al. and <u>U.S. Patent 4,032,661</u>, to Rowsell et al., all of which are herein incorporated by reference. A particularly preferred cooling agent is N-ethyl-p-menthane-3-carboxamide (WS-3 supplied by Sterling Organics), taught by the above incorporated <u>U.S. Patent 4,136,163</u>. Another particularly preferred cooling agent is 3-1-menthoxypropane 1,2-diol (TK-10 supplied by Takasago Perfumery Co., Ltd., Tokyo, Japan). This material is described in detail in <u>U.S. Patent 4,459,425</u>, July 10, 1984 to Amano et al. and incorporated berein by reference.

METHOD OF MANUFACTURE

Good tasting pharmaceutical adsorbate compositions of the present invention are prepared using art-recognized principles and methodologies in mixing the ingredients together and in choosing the type of mixing equipment to be used. However, because of the potential for adverse interactions between the magnesium aluminum silicate and various multiply charged cationic drugs (e.g. chlorpheniramine) - resulting in poor drug dissolution, it is important that certain measures be followed to insure effective drug dissolution. Such a situation is described in McGinty, J.W. and Lach, J.L., In Vitro Adsorption of Various Pharmaceuticals to Montmorillonite, Jour. of Pharm. Sci., 65, 896-902 (1976) and is herein incorporated by reference. By "multiply charged cationic drugs" as used herein, means compounds containing more than one positively charged substituent. The crucial steps involve maintaining the pH in a range equal to or above that of the pKa of the multiply charged cationic drug(s) and, additionally, reserving the addition of said multiply charged cationic drug(s) until after the addition of at least one other cationic compound.

After the ingredients of the present invention are combined, the mixture may then be compounded with effervescent or other water-dispersible substances and dried into dosage forms that rapidly disintegrate upon coming into contact with an aqueous liquid. Suitable effervescent technology is described in chapter 6 of Pharmaceutical Dosage Forms: Tablets, Vol. I, 2nd ed., A Lieberman ed., 1989, Marcel Dekker, Inc. herein incorporated by reference. The above mentioned compounding and drying process may be accomplished by using any of a multitude of solid dosage forming techniques and equipment. Methods of solid dosage formulation are well known in the art and any appropriate method may be utilized. Further information regarding solid dosage formulation can be found in Remington's Pharmaceutical Sciences, pp. 1633-1664, (Alfonso R. Gennaro, editor) (18th ed. 1990).

Alternatively, the resultant fast dissolving dosage form may be achieved by freeze drying. Freeze-drying or lyophilization facilitates disintegration of the composition by forming the dried composition into an open matrix network. In most cases, this results in rapid permeation by the aqueous media, promoting timely delivery of the product's active ingredients. Suitable methods of freeze drying are well known in the art and commonly employed. Any suitable conventional method of freeze-drying or vacuum-drying may be utilized. A preferable method of freezing and drying is to fast freeze the composition and then dry the composition to a final moisture content of about 2% to about 5%. Suitable methods of freeze-drying and production are taught by U.S. Patent 4,642,903, February 17, 1987, to Davies, U.S. Patent 4,946,684, August 7, 1990, to Blank et al., U.S. Patents 4,305,502 and

4.371.516, issued December 15, 1981 and February 1, 1983 respectively, to Gregory et al., and <u>U.S. Patent 5,188,825</u>, February 23,1993, to Hes et al.; which are all incorporated herein by reference

One other form of fast dissolving technology that may be applicable to the present invention is that developed by Janssen Pharmaceutica Inc. and is identified by the trade name QuicksolvTM. This technology is fully described in <u>U.S. Patent 5.215.756</u>, herein incorporated by reference.

EXAMPLES

The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations are possible without departing from the spirit and scope of the invention.

EXAMPLE I

15	Ingredients	W/V%
	chlorpheniramine maleate	0.13300
	phenylpropanolamine HCl	0.83300
	magnesium aluminum silicate ¹	0.50000
	xanthum gum	0.20000
20	potassium sorbate	0.07500
	polysorbate 80	0.10000
	Prosweet™ MM24 ²	0.50000
	sodium saccharin	0.05000
	aspartame	0.30000
25	monoammonium glycyπhizate ³	0.03000
	sucrose	5.00000
	mannitol	10,00000
	3-1-menthoxypropane 1,2-diol ⁴	0.07000
-	N-ethyl-p-menthane-3-carboxamide ⁵	0.02000
30	menthol, natural	0.26600
	peppermint flavor	0.18000
	purified water	q.s. to 100

¹Available as Veegum® HS from R.T. Vanderbilt, Norwalk, CT.

²Available from Virginia Dare Extract Co., Inc., Brooklyn, NY.

³ Available as Magnasweet 185 supplied by McAndrews & Forbes, Camden, NJ.

⁴Available as TK-10 supplied by Takasago Perfumery Co.

⁵Available as WS-3 supplied by Sterling Organics.

In an appropriately sized container, with LightninTM mixer (model #TS2010 (or a high shear homogenizer set at 30 to 50 RPM)) mixing at approximately 250 to 1000 RPM, add the following agents allowing each to dissolve before adding the next: water, phenylpropanolamine HCl, Veegum® HS. Heat, using a hot plate, keeping solution at 77°C, and mix vigorously (250 to 1000 RPM) for 30 minutes. Turn heat off and allow mixture to cool to room temperature (25°C) while mixing for a minimum of an additional 15 minutes. Add monoammonium glycyrrhizate, ProswectTM MM24, potassium sorbate, aspartame, sodium saccharin, WS-3 and peppermint.

Separately, dry blend mannitol, sucrose, and xanthum gum and add slowly to the original mixture. Separately dissolve chlorpheniramine maleate with a sufficient amount of water and add to the original mixture. Adjust the volume to 1.5000 liters and mix for 15 minutes. Separate from the above mixture, in an appropriate container, add polysorbate 80, TK-10, and menthol. Add this mixture to the original mixture while continuing mixing. As the mixing continues, place 1.5 ml of solution into molds and freeze-dry or vacuum-dry the composition. This process, in turn, provides a pharmaceutical adsorbate composition having improved taste. Administration of two tablets to an adult human is the normal and customary dosage for use in treating the symptoms of a respiratory illness or allergy.

20	EXAMPLE II	
	Ingredients	W/V%
	chlorpheniramine maleate	0.13300
•	phenylpropanolamine HCl	0.83300
	magnesium aluminum silicate	0.50000
25	xanthum gum	0.20000
	potassium sorbate	0.07500
	polysorbate 80	0.10000
	Prosweet™ MM24	0.50000
	sodium saccharin	0.05000
30	aspartame	0.30000
	monoammonium glycyrrhizate	0.03000
	sucrose	5.00000
	mannitol	10.00000
	3-1-menthoxypropane 1,2-diol	0.07000
35	N-ethyl-p-menthane-3-carboxamide	0.02000
	menthol, natural	0.26600
	lemon flavor	0.40000

10

•	color, FD&C Yellow #6	0.00240
	purified water	q.s. to 100
	EXAMPLE III	
	Ingredients	<u>W/V%</u>
5	chlorpheniramine maleate	0.13300
	phenylpropanolamine HCl	0.83300
	magnesium aluminum silicate	0.50000
	xanthum gum	0.20000
	potassium sorbate	0.07500
10	polysorbate 80	0.10000
	Prosweet™ MM24	0.50000
	sodium saccharin	0.05000
	aspartame	0.30000
	monoammonium glycyrrhizate	0.03000
15	sucrose	5.00000
	mannitol	10.00000
	3-1-menthoxypropane 1,2-diol	0.07000
	N-ethyl-p-menthane-3-carboxamide	0.02000
	menthol, natural	0.26600
20	raspberry flavor	0.30000
	color, FD&C Yellow #6	0.00236
	color, FD&C Red #33	0.00326
	purified water	q.s. to 100
	EXAMPLE IV	
25	Ingredients	<u>W/V%</u>
	diphenhydramine HCl	0.83300
	phenylpropanolamine HCl	0.83300
•	magnesium aluminum silicate	0.50000
	xanthum gum	0.20000
30	potassium sorbate	0.07500
	polysorbate 80	0.10000
	Prosweet™ MM24	0.50000
	sodium saccharin	0.15000
	aspartame	0.30000
35	monoammonium glycyrrhizate	0.03000
	sucrose	5.00000
•	mannitol	10.00000

	3-1-menthoxypropane 1,2-diol	0.07000
	N-ethyl-p-menthane-3-carboxamide	0.02000
	menthol, natural	0.26600
	lemon flavor	0.20000
, 5	menthol monophosphate	0.30000
	purified water	q.s. to 100
	EXAMPLE V	•
	Ingredients	<u>W/V%</u>
	triprolidine HCI	0.16700
10	phenylpropanolamine HCl	0.83300
	magnesium aluminum silicate	0.50000
	ascorbic acid	2.00000
	xanthum gum	0.20000
	potassium sorbate	0.07500
15	polysorbate 80	0.10000
	Prosweet™ MM24	1.00000
	sodium saccharin	0.15000
	aspartame	1.00000
	monoammonium glycyrrhizate	1.00000
.20	sucrose	5.00000
	mannitol	10.00000
	3-1-menthoxypropane 1,2-diol	0.07000
	N-ethyl-p-menthane-3-carboxamide	0.02000
	menthol, natural	0.26600
25	lemon flavor	0.40000
	purified water	q.s. to 100
	EXAMPLE VI	
	Ingredients	<u>W/V%</u>
	chlorpheniramine maleate	0.13300
30	phenylpropanolamine HCl	0.83300
	magnesium aluminum silicate	2.00000
	calcium carbonate l	0.80000
	xanthum gum	0.20000
	potassium sorbate	0.07500
35	polysorbate 80	0.10000
	Prosweet™ MM24	0.50000
	sodium saccharin	0.15000

	purified water	q.s. to 100
	peppermint flavor	0.18000
	menthol, natural	0.26600
	N-ethyl-p-menthane-3-carboxamide	0.02000
5	3-1-menthoxypropane 1,2-diol	0.07000
	mannitol	10.00000
	sucrose	5.00000
	monoammonium glycyrrhizate	0.03000
	aspartame	0.30000

10

15

¹Granulated calcium carbonate containing 93.3% calcium carbonate, 6.3% glucose and 0.4% gelatin; supplied by Whittaker, Clark & Daniels, Philadelphia, PA.

Examples II-VI are further examples of combinations used in treating the symptoms of a respiratory illness or allergy in a human and are manufactured in a manner substantially similar to Example I. Administration of two tablets is the normal and customary dosage.

EXAMPLE VII

	<u>Ingredients</u>	<u>W/V%</u>
	magnesium aluminum silicate	8,48900
	loperamide	2.44000
20	polysorbate 80	0.04000
	sodium saccharin	0.05200
	aspartame	0.05200
	sucrose	14,00000
	mannitol	12.25500
25	Novagel RCN-151	1,60000
	lemon Flavor	0.47200
	vanilla Creme	0.15000
	citric acid	0.45000
	purified water	q.s. to 100
30	EXAMPLE VIII	
	<u>Ingredients</u>	<u>W/V%</u>
	magnesium aluminum silicate	4.00000
	bismuth subsalicylate	21.69930
	polysorbate 80	0.03500
35	sodium saccharin	0.04550
	aspartame	1.00000
	sucrose	2.00000

	mannitol	4.15000
	Novagel RCN-151	1.40000
	strawberry flavor	0.54600
	vanilla creme	0.54990
5	citric acid	0.06690
	Klucel EF NF ²	0.25200
	Antifoam AF3	0.21000
	purified water	q.s. to 100
	EXAMPLE D	<u>K</u>
10	Ingredients	W/V%
	magnesium aluminum silicate	4.00000
	simethicone	3.50000
	polysorbate 80	0.03500
	sodium saccharin	0.04550
15	aspartame	0.04550
	sorbitol	0.70000
	sucrose	8.25000
	mannitol	16.29800
•	Novagel RCN-151	1.40000
20	cherry flavor	0.19460
	vanilla creme	0.24780
	citric acid	0.28320
	Antifoam AF ³	0.21000
	purified water	q.s. to 100

25 ^ISupplied by FMC Corporation, Philadelphia, Pennsylvania

30

Examples VII -IX are further examples of combinations used in treating the symptoms of gastrointestinal illnesses in humans and are manufactured in a manner substantially similar to Example I. Administration of two tablets is the normal and customary dosage.

²Supplied by AQUALON, Hopewell, Virginia

³Supplied by Dow Corning, Midland, Michigan

What is Claimed is:

- A pharmaceutical composition comprising:
 - (a) a clay component consisting of magnesium aluminum silicate:
 - (b) a safe and effective amount of at least two pharmaceutically acceptable active ingredients selected from the group consisting of: an antitussive; an antinauseant; a nutritional supplement: a laxative; an appetite suppressant; an analgesic; an antiasthmatic; an antihistamine; a decongestant; an expectorant; an antacid; an antidiarrheal, a H₂-Receptor antagonist and mixtures thereof, preferably phenylpropanolamine and chlorpheniramine; and
 - (c) a pharmaceutically acceptable carrier wherein said pharmaceutically acceptable carrier can be rapidly disintegrated in aqueous solution.
- A pharmaceutical composition according to Claim 1 wherein said carrier is a freeze-dried matrix, an effervescent system, or a liquid/liquid extract system.
- 3. A pharmaceutical composition according to any one of the preceding Claims wherein the composition additionally contains one or more sweetening agents, preferably selected from the group consisting of sodium saccharin, aspartame, monoammonium glycyrrhizate, sucrose, mannitol and mixtures thereof.
- 4. A pharmaceutical composition according to any one of the preceding Claims wherein the composition additionally contains one or more flavoring agents.
- A pharmaceutical composition according to any one of the preceding Claims wherein the composition additionally contains one or more releasing agents.
- 6. A pharmaceutical composition according to any one of the preceding Claims wherein the composition additionally contains one or more cooling agents.
- A pharmaceutical composition according to any one of the preceding Claims additionally comprising a flavoring agent selected from the group consisting of: menthol, peppermint, spearmint, raspberry, cherry, orange, vanilla, anise, blueberry, banana, chocolate, caramel, strawberry, lemon, grape and mixtures thereof.

- 8. A pharmaceutical composition according to any one of the preceding Claims additionally comprising a releasing agent selected from the group consisting of: polysorbate 80, sodium lauryl sulfate, magnesium stearate and mixtures thereof.
- 9. A pharmaceutical composition according to any one of the preceding Claims additionally comprising a cooling agent selected from the group consisting of 3-1-methoxypropane 1,2-diol, N-ethyl-p-methane-3-carboxamide and mixtures thereof.
- 10. A pharmaceutical composition according to any one of the preceding Claims wherein the ratio of said magnesium aluminum silicate to said phenylpropanolamine and said chlorpheniramine is 1 to 5.0 and 1 to 0.8 respectively.

cited in search report	date	member(s)		date
WO-A-9217164	15-10-92	AU-A-	1761492	02-11-92
		BR-A-	9205827	28-06-94
		CZ-A-	9302260	13-04-94
		EP-A-	0578768	19-01-94
		HU-A-	65881	28-07-94
		JP-T-	6506682	28-07-94
		US-A-	5244670	14-09-93
EP-A-0239542	30-09-87	US-A-	4711774	08-12-87
		AU-B-	574605	07-07-88
		AU-A-	7069187	01-10-87
		CA-A-	1282328	02-04-91
		JP-B-	6076316	28-09-94
		JP-A-	62238215	19-10-87
		us-a-	4758424	19-07-88
	•	US-A-	4758425	19-07-88
		US-A-	4761274	02-08-88
		ZA-A-	8701660	27-08-87
WO-A-9420074	15-09-94	NONE		